

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 December 2000 (07.12.2000)

PCT


(10) International Publication Number  
**WO 00/72812 A1**

- (51) International Patent Classification<sup>7</sup>: A61K 7/06, Chester, OH 45069 (US). YOUNGQUIST, Robert, Scott [US/US]; 8511 Charleston Knoll Court, Mason, OH 45040 (US).  
A61P 17/14
- (21) International Application Number: PCT/US00/05253
- (22) International Filing Date: 1 March 2000 (01.03.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/137,022 1 June 1999 (01.06.1999) US
- (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).
- (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).
- (81) Designated States (national): AU, BR, CA, CN, JP, KR, MX, US.
- (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- Published:  
— With international search report.
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): ZHANG, Lilly, Li-Xin [US/US]; 6537 Meadowbrook Court, West

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF TREATING HAIR LOSS USING DIPHENYLETHER DERIVATIVES

WO 00/72812 A1



(I)

(57) Abstract: The present disclosure describes methods for treating hair loss in mammals, including arresting and/or reversing hair loss and promoting hair growth. The methods comprise administering a cardiac-sparing compound having a structure of formula (I) as described herein and a pharmaceutically-acceptable carrier.

## METHOD OF TREATING HAIR LOSS USING DIPHENYLETHER DERIVATIVES

### FIELD OF THE INVENTION

The present invention relates to methods for treating hair loss in mammals, including arresting and / or reversing hair loss and promoting hair growth.

### BACKGROUND OF THE INVENTION

Hair loss is a common problem which occurs, for example, through natural processes or is often chemically promoted through the use of certain therapeutic drugs designed to alleviate conditions such as cancer. Often such hair loss is accompanied by lack of hair regrowth which causes partial or full baldness.

As is well-known in the art, hair growth occurs by a cycle of activity which involves alternating periods of growth and rest. This cycle is often divided into three main stages which are known as anagen, catagen, and telogen. Anagen is the growth phase of the cycle and may be characterized by penetration of the hair follicle deep into the dermis with rapid proliferation of cells which are differentiating to form hair. The next phase is catagen, which is a transitional stage marked by the cessation of cell division, and during which the hair follicle regresses through the dermis and hair growth is ceased. The next phase, telogen, is often characterized as the resting stage during which the regressed follicle contains a germ with tightly packed dermal papilla cells. At telogen, the initiation of a new anagen phase is caused by rapid cell proliferation in the germ, expansion of the dermal papilla, and elaboration of basement membrane components. Wherein hair growth ceases, most of the hair follicles reside in telogen and anagen is not engaged, thus causing the onset of full or partial baldness.

There have been many attempts in the literature to invoke the regrowth of hair by, for example, the promotion or prolongation of anagen. Currently, there are two drugs approved by the United States Food and Drug Administration for the treatment of male pattern baldness: topical minoxidil (marketed as Rogaine® by Pharmacia & Upjohn), and oral finasteride (marketed as Propecia® by Merck & Co., Inc.). For several reasons, however, including safety concerns and / or lack of efficacy, the search for efficacious hair growth inducers is ongoing.


Interestingly, it is known that the thyroid hormone known as thyroxine ("T4") converts to thyronine ("T3") in human skin by deiodinase I, a selenoprotein. Selenium deficiency causes a decrease in T3 levels due to a decrease in deiodinase I activity; this

reduction in T3 levels is strongly associated with hair loss. Consistent with this observation, hair growth is a reported side effect of administration of T4. See, e.g., Berman, "Peripheral Effects of L-Thyroxine on Hair Growth and Coloration in Cattle", *Journal of Endocrinology*, Vol. 20, pp. 282 - 292 (1960); and Gunaratnam, "The Effects of Thyroxine on Hair Growth in the Dog", *J. Small Anim. Pract.*, Vol. 27, pp. 17 - 29 (1986). Furthermore, T3 and T4 have been the subject of several patent publications relating to treatment of hair loss. See, e.g., Fischer et al., DE 1,617,477, published January 8, 1970; Mortimer, GB 2,138,286, published October 24, 1984; and Lindenbaum, WO 96/25943, assigned to Life Medical Sciences, Inc., published August 29, 1996.

Unfortunately, however, administration of T3 and / or T4 to treat hair loss is not practicable because these thyroid hormones are also known to induce significant cardiotoxicity. See, e.g., Walker et al., U.S. Patent No. 5,284,971, assigned to Syntex, issued February 8, 1994 and Emmett et al., U.S. Patent No. 5,061,798, assigned to Smith Kline & French Laboratories, issued October 29, 1991. Surprisingly, however, the present inventors have discovered biphenyl derivatives which promote hair growth without inducing cardiotoxicity. Consistent with this discovery, but without intending to be limited by theory, the present inventors have surprisingly discovered that the biphenyl derivatives useful in the present invention interact strongly with hair-selective thyroid hormone receptors but interact less strongly, or not at all, with heart-selective hormone receptors. These unique properties are, of course, not shared with T3 and / or T4. Accordingly, the biphenyl derivatives described for use in the methods and compositions herein are cardiac-sparing compounds useful for treating hair loss, including arresting and / or reversing hair loss and promoting hair growth.

#### SUMMARY OF THE INVENTION

The present invention relates to methods for treating hair loss comprising administering a cardiac-sparing compound which has been found by the present inventors to be particularly useful for treating hair loss in mammals, including arresting and / or reversing hair loss and promoting hair growth. The compounds utilized in the present method are biphenyl derivatives having the structure:



and pharmaceutically acceptable salts, hydrates, and biohydrolyzable amides, esters, and imides thereof, wherein R, R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, X, W, and Z are defined herein.

### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to methods of using compounds and compositions which are particularly useful for treating hair loss in mammals, including arresting and / or reversing hair loss and promoting hair growth.

In addition to discovering that the present compounds are useful for treating hair loss, the present inventors have also surprisingly discovered that the preferred compounds are cardiac-sparing. The preferred compounds useful in the method of the present invention are therefore, as defined herein below, cardiac-sparing.

Publications and patents are referred to throughout this disclosure. All references cited herein are hereby incorporated by reference.

All percentages, ratios, and proportions used herein are by weight unless otherwise specified.

In the description of the invention various embodiments and/or individual features are disclosed. As will be apparent to the ordinarily skilled practitioner all combinations of such embodiments and features are possible and can result in preferred executions of the invention.

As used herein, wherein any variable, moiety, group, or the like occurs more than one time in any variable or structure, its definition at each occurrence is independent of its definition at every other occurrence.

### Definition and Usage of Terms

The following is a list of definitions for terms used herein:

As used herein "salt" is a cationic salt formed at any acidic (*e.g.*, carboxyl) group, or an anionic salt formed at any basic (*e.g.*, amino) group. Many such salts are known in the art.

Preferred cationic salts include the alkali metal salts (such as, for example, sodium and potassium), alkaline earth metal salts (such as, for example, magnesium and calcium), and organic salts. Preferred anionic salts include the halides (such as, for example, chloride salts). Such acceptable salts must, when administered, be appropriate for mammalian use.

As used herein, "alkyl" is a saturated, straight or branched chain monovalent hydrocarbon radical. Unless otherwise specified, alkyls have from 1 to about 10 carbon atoms ( $C_1 - C_{10}$ ). Preferred alkyls include, for example, methyl, ethyl, propyl, *iso*-propyl, *tert*-butyl, *n*-butyl, *sec*-butyl, and *iso*-butyl.

As used herein, the term "aryl" is a carbocyclic or heterocyclic aryl. Carbocyclic aryl is optionally substituted phenyl or optionally substituted naphthyl. Heterocyclic aryl is optionally substituted phenyl or optionally substituted naphthyl, having at least one heteroatom (N, O, or S) making up the aryl ring.

As used herein, the term "aryl-lower alkyl" is aryl substituted by at least one alkyl, lower alkoxy, lower alkanoyloxy, or trifluoromethyl.

As used herein, "biohydrolyzable amides" are amides of the compounds used in the present invention which do not interfere with the activity of the compound, or that are readily converted *in vivo* by a mammalian subject to yield an active compound.

As used herein, "biohydrolyzable esters" are esters of the compounds used in the present invention which do not interfere with the activity of the compound, or that are readily converted *in vivo* by a mammalian subject to yield an active compound.

As used herein, "biohydrolyzable imides" are imides of the compounds used in the present invention which do not interfere with the activity of the compound, or that are readily converted *in vivo* by a mammalian subject to yield an active compound.

As used herein "cycloalkyl" refers to a saturated cyclic hydrocarbon radical, preferably  $C_5 - C_7$  cycloalkyl. Most preferably, a cycloalkyl is cyclopentyl or cyclohexyl.

As used herein, the term "cycloalkyl-lower alkyl" is cycloalkyl substituted by at least one alkyl, lower alkoxy, lower alkanoyloxy, or trifluoromethyl. Non-limiting examples of cycloalkyl-lower alkyl include 1- or 2-(cyclopentyl or cyclohexyl)ethyl; 1-, 2-, or 3-(cyclopentyl or cyclohexyl)propyl; or 1-, 2-, 3-, or 4-(cyclopentyl or cyclohexyl)butyl.

As used herein "halogen", "halo", or the like refers to chlorine, bromine, iodine, and fluorine, preferably fluorine and chlorine.

As used herein, the term "lower" in connection with organic radicals or compounds defines such with up to and including 7, preferably up to and including 4, and most preferably one or two carbon atoms. The "lower" radical or compound may be straight or branched.

As used herein, "pharmaceutically acceptable" means suitable for use in a human or other mammal.

As used herein, "safe and effective amount of a compound" (or composition, or the like) means an amount that is effective to exhibit biological activity, preferably wherein the biological activity is arresting and / or reversing hair loss or promoting hair growth, at the site(s) of activity in a mammalian subject, without undue adverse side effects (such as toxicity, irritation, or allergic response), commensurate with a reasonable benefit / risk ratio when used in the manner of this invention.


Acyl is preferably lower alkanoyl, carbocyclic aryl-lower alkanoyl, or carbocyclic aroyl.

Lower alkanoyl is preferably acetyl, propionyl, butyryl, or pivaloyl.

Lower alkanoyloxy is preferably acetoxy, pivaloyloxy, or propionyloxy.


#### Methods of the Present Invention

The present invention relates to methods of treating hair loss comprising administering a composition comprising a compound having the structure:



and pharmaceutically acceptable salts, hydrates, and biohydrolyzable amides, esters, and imides thereof, wherein:

- (a) R is selected from the group consisting of hydrogen, hydroxy, esterified hydroxy, and etherified hydroxy;
- (b) R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> are each, independently, selected from the group consisting of hydrogen, halogen, trifluoromethyl, and lower alkyl;
- (c) R<sub>3</sub> is selected from the group consisting of halogen, trifluoromethyl, lower alkyl, aryl, aryl-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, and:



- (d)  $R_8$  is selected from the group consisting of hydrogen, lower alkyl, aryl, cycloalkyl, aryl-lower alkyl, and cycloalkyl-lower alkyl;
- (e)  $R_9$  is selected from the group consisting of hydroxy and acyloxy;
- (f)  $R_{10}$  is selected from the group consisting of hydrogen and lower alkyl; or wherein  $R_9$  and  $R_{10}$  together represent oxo;
- (g) W is selected from the group consisting of -O- and -S-;
- (h) X is selected from the group consisting of -NR<sub>7</sub>, S, and O;
- (i)  $R_5$  is selected from the group consisting of hydrogen, lower alkyl, and aryl-lower alkyl and  $R_6$  is hydrogen; or wherein  $R_5$  and  $R_6$  are together oxo, provided that X is -NR<sub>7</sub>;
- (j)  $R_7$  is selected from the group consisting of hydrogen and lower alkyl; and
- (k) Z is selected from the group consisting of carboxyl and carboxyl derivatized as a pharmaceutically acceptable ester or amide.

The compounds useful in the method herein are further described in Yokoyama et al., U.S. Patent No. 5,401,772, assigned to Ciba-Geigy Corp., issued March 28, 1995; Yokoyama et al., EP 0,580,550, assigned to Ciba-Geigy Corp., published January 26, 1994; and Yokoyama et al., "Synthesis and Structure-Activity Relationships of Oxamic Acid and Acetic Acid Derivative Related to L-Thyronine", *Journal of Medicinal Chemistry*, Vol. 38, pp. 695 - 707 (1995). However, for convenience, the compounds are more fully described herein below:

#### The R Moiety

The R moiety is selected from hydrogen, hydroxy, esterified hydroxy, and etherified hydroxy. As used herein "esterified hydroxy" refers to acyloxy, *e.g.*, acyloxy derived from an organic carboxylic acid. Preferred esterified hydroxy include lower alkanoyloxy, aroyloxy, and aryl-lower alkanoyloxy. As used herein, "etherified hydroxy" preferably represents lower alkoxy, lower alkenyloxy, C<sub>5</sub> - C<sub>7</sub> cycloalkyloxy, carbocyclic aryl-lower alkoxy, tetrahydropyranyloxy, C<sub>5</sub> - C<sub>7</sub> cycloalkyl-lower alkoxy, and the like.

More preferably, the R moiety is selected from hydroxy, esterified hydroxy, and etherified hydroxy. Even more preferably, the R moiety is selected from hydroxy, lower

alkanoyloxy, lower alkoxy, and tetrahydropyranyloxy. Most preferably, the R moiety is hydroxy.

Preferably, the R moiety substitutes at the 4' position as shown herein.

#### The R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> Moieties

R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> are each, independently, selected from hydrogen, halogen, trifluoromethyl and lower alkyl.


Preferably, R<sub>1</sub> and R<sub>2</sub> are each, independently, selected from halogen, trifluoromethyl, and C<sub>1</sub> - C<sub>3</sub> alkyl. More preferably, R<sub>1</sub> and R<sub>2</sub> are each, independently, selected from halogen and C<sub>1</sub> - C<sub>3</sub> alkyl. Still more preferably, R<sub>1</sub> and R<sub>2</sub> are each, independently, selected from chlorine and methyl. Preferably, R<sub>1</sub> and R<sub>2</sub> are equivalent. Most preferably, R<sub>1</sub> and R<sub>2</sub> are each methyl.

R<sub>4</sub> is preferably hydrogen.

Preferably, R<sub>1</sub> substitutes at the 3 position as shown herein. Preferably, R<sub>2</sub> substitutes at the 5 position as shown herein. Preferably, R<sub>4</sub> substitutes at the 5' position as shown herein.


#### The R<sub>3</sub> Moiety

The R<sub>3</sub> moiety is selected from halogen, trifluoromethyl, lower alkyl, aryl, aryl-lower alkyl, cycloalkyl-lower alkyl, and:



R<sub>8</sub> is selected from hydrogen, lower alkyl, aryl, cycloalkyl, aryl-lower alkyl, and cycloalkyl-lower alkyl. R<sub>9</sub> is selected from hydroxy and acyloxy. As used herein, the term "acyloxy" is -O-acyl, wherein acyl is preferably selected from lower alkanoyl, carbocyclic aryl-lower alkanoyl, and carbocyclic aroyl. R<sub>10</sub> is selected from hydrogen and lower alkyl; or wherein R<sub>9</sub> and R<sub>10</sub> together represent oxo (doubly-bonded oxygen).

Preferably, the R<sub>3</sub> moiety is selected from lower alkyl, aryl-lower alkyl, cycloalkyl-lower alkyl and





More preferably, the R<sub>3</sub> moiety is selected from *iso*-propyl, benzyl, benzyl substituted with halogen, and trifluoromethyl. Most preferably, R<sub>3</sub> is *iso*-propyl.

The R<sub>3</sub> moiety preferably substitutes at the 3' position as shown herein.

#### The W Moiety

W is selected from -O- and -S-. Most preferably, W is -O-.

#### The X Moiety

X is selected from -NR<sub>7</sub>-, -S-, and -O-. Preferably, X is -NR<sub>7</sub>- or -O-. Most preferably, X is -NR<sub>7</sub>-.

#### The R<sub>5</sub> and R<sub>6</sub> Moieties

R<sub>5</sub> is selected from hydrogen, lower alkyl, and aryl-lower alkyl. R<sub>6</sub> is hydrogen. Alternatively, and most preferably, R<sub>5</sub> and R<sub>6</sub> are, together, an oxo moiety (=O).

#### The R<sub>7</sub> Moiety

R<sub>7</sub> is selected from hydrogen and lower alkyl. Most preferably, R<sub>7</sub> is hydrogen.

#### The Z Moiety

Z is selected from carboxyl (-CO<sub>2</sub>H) and carboxyl derivatized as a pharmaceutically acceptable ester or amide. As used herein, a "carboxyl derivatized as a pharmaceutically acceptable ester" represents esterified carboxyl, preferably a prodrug ester which is convertible by solvolysis or under physiological conditions to the corresponding free carboxylic acid. Similarly, a "carboxyl derivatized as a pharmaceutically acceptable amide" represents carboxyl which has been functionalized as an amide, preferably a prodrug amide which is convertible by solvolysis or under physiological conditions to the corresponding free carboxylic acid. The amide may be a primary amide, *i.e.*, -C(=O)NH<sub>2</sub>.

Preferably, Z is selected from carboxyl and carboxyl derivatized as a pharmaceutically acceptable ester. Most preferably, Z is carboxyl.

Preferred carboxyl derivatized as a pharmaceutically acceptable ester include lower alkoxy-carbonyl; (amino, acylamino, mono- or di-lower alkylamino)-lower alkoxy-carbonyl; carboxy-lower alkoxy-carbonyl, *e.g.*,  $\alpha$ -carboxy-lower alkoxy-carbonyl; lower alkoxy-carbonyl-lower alkoxy-carbonyl, *e.g.*,  $\alpha$ -lower alkoxy-carbonyl-lower alkoxy-carbonyl;  $\alpha$ -(di-lower alkylamino, amino, mono-lower alkylamino, morpholino, piperidino, pyrrolidino, 1-lower


alkyl-piperazino)-carbonyl-lower alkoxy carbonyl; carbocyclic and heterocyclic aryl-lower alkoxy carbonyl, preferably optionally (halo, lower alkyl or lower alkoxy)-substituted benzyloxy carbonyl, or pyridylmethoxy carbonyl; 1-(hydroxy, lower alkanoyloxy or lower alkoxy)-lower alkoxy carbonyl, *e.g.*, pivaloyloxymethoxy carbonyl; (hydroxy, lower alkanoyloxy or lower alkoxy)-lower alkoxy methoxy carbonyl; 1-(lower alkoxy carbonyloxy)-lower alkoxy carbonyl; 5-indanyloxy carbonyl; 3-phthalidoxycarbonyl and (lower alkyl, lower alkoxy or halo)-substituted 3-phthalidoxycarbonyl; dihydroxypropyloxy carbonyl wherein hydroxy groups are free or are protected in the form of ketals, *e.g.*, a lower alkylidene, a benzylidene or a 5- or 6-membered cycloalkylidene derivative, preferably (2,2-dimethyl-1,3-dioxolan-4-yl)-methoxy carbonyl.

Carboxyl derivatized as a pharmaceutically acceptable prodrug ester is most preferably C<sub>1</sub> - C<sub>4</sub> alkoxy carbonyl, benzyloxy carbonyl optionally substituted on phenyl by lower alkyl, lower alkoxy, halo or trifluoromethyl, 1-(C<sub>2</sub> - C<sub>4</sub>-alkanoyloxy)-ethoxy carbonyl, (2,2-dimethyl-1,3-dioxolan-4-yl)-methoxy carbonyl, 5-indanyloxy carbonyl, 1-(C<sub>1</sub> - C<sub>4</sub>-alkoxy carbonyloxy)-ethoxy carbonyl or 3-pyridylmethoxy carbonyl.

Carboxyl derivatized as a pharmaceutically acceptable amide is preferably carbamoyl or N-substituted carbamoyl, preferably lower alkylamino carbamoyl, arylamino carbamoyl, di-lower alkylamino carbamoyl, morpholino carbamoyl, N-lower alkylpiperazino carbamoyl, pyrrolidino carbamoyl, piperidino carbamoyl, (amino or acylamino)-lower alkylamino carbamoyl or aryl-lower alkylamino carbamoyl.


#### Preferred Compounds of the Present Invention

Preferred compounds of the present invention include those wherein X is -NR<sub>7</sub> and R<sub>5</sub> and R<sub>6</sub> are together oxo. Such compounds are represented as:



An even more preferred compound of this structure is wherein W is -O-.


Further preferred compounds include those wherein R substitutes at the 4' position, R<sub>3</sub> substitutes at the 3' position, R<sub>1</sub> substitutes at the 3 position, R<sub>2</sub> substitutes at the 5 position, and R<sub>4</sub> is hydrogen. Such compounds are represented as:



An even more preferred compound of this structure is wherein W is -O-.

Other preferred compounds useful in the present invention are described in Yokoyama et al., "Synthesis and Structure - Activity Relationships of Oxamic Acid and Acetic Acid Derivatives Related to L-Thyronine", *Journal of Medicinal Chemistry*, Vol. 38, pp. 695 - 707 (1995). These compounds are further described in Table 1 below:

Table 1



Example No.	R <sub>1</sub>	R <sub>2</sub>	Y	R <sub>7</sub>
1	hydrogen	hydrogen	hydroxy	hydrogen
2	iodine	iodine	hydroxy	hydrogen
3	bromine	bromine	hydroxy	hydrogen
4	bromine	bromine	NH <sub>2</sub>	hydrogen
5	bromine	bromine	NH-CH <sub>3</sub>	hydrogen
6	chlorine	chlorine	hydroxy	hydrogen

7	fluorine	fluorine	hydroxy	hydrogen
8	methyl	methyl	hydroxy	hydrogen
9	methyl	methyl	NH <sub>2</sub>	hydrogen
10	methyl	methyl	hydroxy	methyl
11	iso-propyl	iso-propyl	hydroxy	hydrogen
12	methyl	methyl	O-CH <sub>2</sub> -CH <sub>3</sub>	hydrogen
13	methyl	methyl	O-CH <sub>2</sub> -phenyl	hydrogen

### Analytical Methods

The present invention relates to methods of treating hair loss by administering a compound having a structure as described herein. Preferably, the compound utilized in the present invention will be cardiac-sparing. Compounds (test compounds) may be tested for their ability to induce anagen and their lack of cardiotoxicity (cardiac-sparing) using the following methods. Alternatively, other methods well-known in the art may be used (but with the term "cardiac-sparing" being defined according to the method disclosed herein below).

### Cardiotoxicity Assay:

The cardiotoxicity assay measures the potential of a test compound to adversely affect the cardiovascular system. As thyroid hormone (T3) damages the cardiovascular system, the heart enlarges. See, e.g., Gomberg-Maitland et al., "Thyroid hormone and Cardiovascular Disease", *American Heart Journal*, Vol. 135(2), pp. 187-196 (1998); Klein and Ojamaa, "Thyroid Hormone and the Cardiovascular System", *Current Opinion in Endocrinology and Diabetes*, Vol. 4, pp.341-346 (1997); and Klemperer et al., "Thyroid Hormone Therapy and Cardiovascular Disease", *Progress in Cardiovascular Diseases*, Vol. 37 (4), pp. 329-336 (1996). This increases the weight of the heart relative to whole body weight. The cardiotoxicity assay herein below is used to test compounds for potentially adverse cardiac effects by measuring their effect on the heart-to-body weight ratio.

Two groups each of six male Sprague Dawley rats (Harlan Sprague Dawley, Inc., Indianapolis, IN) (each weighing from approximately 220 grams to 235 grams) are utilized. The first group is a vehicle control group and the second group is a test compound group. The length of the assay is 30 days, with treatment of vehicle or test compound in vehicle daily for 28 of those days as described below.

Prior to initiation of the assay, each rat is allowed to acclimate to standard environmental conditions for 5 days. Each rat receives food (standard rat chow diet) and water *ad libitum* 5 days prior to initiation of the assay as well as to termination of the study.

The vehicle is 91:9 (v:v) propylene glycol:ethanol. The test compound is prepared at a concentration of 500 µg/mL in the vehicle.

Each rat is weighed on day 1 of the assay. Dosage calculations are then performed: each rat will be administered daily a dosing solution of vehicle or test compound in vehicle (depending on whether the rat is in the vehicle control group or the test compound group, respectively) at 500 µL of dosing solution per kg of rat. For rats in the test compound group, this corresponds to a dose of 250 µg of test compound per kg of rat.

Day 2 is the first day of treatment with dosing solution for both groups. Body weights are taken for each rat on days 3, 5, 8, 10, 12, 15, 17, 19, 22, 24, 26, and 29 prior to dosing for that day; for each rat, the dosing solutions are recalculated and administered accordingly upon change in body weight.

Treatment occurs once daily in the morning on days 2 through 29, inclusive, for each rat in each group. For each treatment, the dosing solution is administered subcutaneously between the shoulders of the rat such that the injection sites are rotated in this area.

On day 30 in the morning, the rats of each group are euthanized with CO<sub>2</sub> from dry ice. Each rat is immediately weighed for total body weight.

The hearts of each rat are then excised as follows. An incision is made to expose the abdominal cavity. The rib cage is carefully cut at the sternum with small scissors, such that the heart and lungs are exposed. With small scissors and forceps, the vessels connected to the heart are cut away from the heart. These vessels include the caudal vena cava, left cranial vena cava (pulmonary trunk), right cranial vena cava, thoracic aorta, right subclavian artery, internal thoracic artery and vein, and any other small attachments. The heart is then immediately taken out intact, including the left and right auricles and left and right ventricles. Immediately thereafter, any excess tissue is trimmed away, the heart is lightly blotted on a paper towel until no more blood is visibly left behind on the paper towel, and the heart is weighed.

The heart weight is divided by the body weight after euthanization for each rat to give the heart/body ratio. The heart/body ratios for each rat in the vehicle control group are added together and divided by 6 (*i.e.*, the total number of rats in the group) to give RV (ratio for vehicle control group). Similarly, the heart/body ratios for each rat in the test compound group are added together and divided by 6 to give RT (ratio for test compound group).

The index C is then calculated by dividing RT by RV. As defined herein, where C is less than 1.3, the test compound is cardiac-sparing. Preferably, C is less than 1.2, more preferably less than 1.15, and most preferably less than 1.1. In accordance with this method, T3 and T4 are not cardiac-sparing.

#### Telogen Conversion Assay:

The Telogen Conversion Assay measures the potential of a test compound to convert mice in the resting stage of the hair growth cycle ("telogen"), to the growth stage of the hair growth cycle ("anagen").

Without intending to be limited by theory, there are three principal phases of the hair growth cycle: anagen, catagen, and telogen. It is believed that there is a longer telogen period in C3H mice (Harlan Sprague Dawley, Inc., Indianapolis, IN) from approximately 40 days of age until about 75 days of age, when hair growth is synchronized. It is believed that after 75 days of age, hair growth is no longer synchronized. Wherein about 40 day-old mice with dark fur (brown or black) are used in hair growth experiments, melanogenesis occurs along with hair (fur) growth wherein the topical application of hair growth inducers are evaluated. The Telogen Conversion Assay herein below is used to screen compounds for potential hair growth by measuring melanogenesis.

Three groups of 44 day-old C3H mice are utilized: a vehicle control group, a positive control group, and a test compound group, wherein the test compound group is administered a compound used in the method of the present invention. The length of the assay is at least 19 days with 15 treatment days (wherein the treatment days occur Mondays through Fridays). Day 1 is the first day of treatment. Most studies will end on Day 19, but a few may be carried out to Day 24 if the melanogenesis response looks positive, but occurs slowly. A typical study design is shown in Table 1 below. Typical dosage concentrations are set forth in Table 1, however the skilled artisan will readily understand that such concentrations may be modified.

Table 1

Group #	Animal #	Compound	Concentration	Application volume	Length of Study
1	1 - 10	Test Compound	0.1% in vehicle*	400 $\mu$ L topical	19 or 24 days

			*		
2	11 - 20	Positive Control (T3)	0.01% in vehicle* *	400 $\mu$ L topical	19 or 24 days
3	21 - 30	Vehicle**	N/A	400 $\mu$ L topical	19 or 24 days

\*\*The vehicle is 60% ethanol, 20% propylene glycol, and 20% dimethyl isosorbide (commercially available from Sigma Chemical Co., St. Louis, MO).

The mice are treated topically Monday through Friday on their lower back (base of tail to the lower rib). A pipettor and tip are used to deliver 400  $\mu$ L to each mouse's back. The 400  $\mu$ L application is applied slowly while moving hair on the mouse to allow the application to reach the skin.

While each treatment is being applied to the mouse topically, a visual grade of from 0 to 4 will be given to the skin color in the application area of each animal. As a mouse converts from telogen to anagen, its skin color will become more bluish-black. As indicated in Table 2, the grades 0 to 4 represent the following visual observations as the skin progresses from white to bluish-black.

Table 2

<u>Visual Observation</u>	<u>Grade</u>
Whitish Skin Color	0
Skin is light gray (indication of initiation of anagen)	1
Appearance of Blue Spots	2
Blue Spots are aggregating to form one large blue area	3
Skin is dark blue (almost black) with color covering majority of treatment area (indication of mouse in full anagen)	4

#### Method of Making

The compounds used in the methods of the present invention are prepared according to procedures which are well-known to those ordinarily skilled in the art. The starting materials used in preparing the compounds are known, made by known methods, or are commercially available as a starting material.

It is recognized that the ordinarily skilled artisan in the art of organic chemistry can readily carry out standard manipulations of organic compounds without further direction.

Examples of such manipulations are discussed in standard texts such as J. March, Advanced Organic Chemistry, John Wiley & Sons (1992).

The ordinarily skilled artisan will readily appreciate that certain reactions are best carried out when other functionalities are masked or protected in the compound, thus increasing the yield of the reaction and / or avoiding any undesirable side reactions. Often, the artisan utilizes protecting groups to accomplish such increased yields or to avoid the undesired reactions. These reactions are found in the literature and are also well within the scope of the skilled artisan. Examples of many such manipulations can be found in, for example, T. Greene, Protecting Groups in Organic Synthesis, John Wiley & Sons (1981).


The compounds of the present invention may have one or more chiral centers. As a result, one may selectively prepare one optical isomer, including diastereomers and enantiomers, over another, for example by chiral starting materials, catalysts or solvents, or may prepare both stereoisomers or both optical isomers, including diastereomers and enantiomers at once (a racemic mixture). Since the compounds of the invention may exist as racemic mixtures, mixtures of optical isomers, including diastereomers and enantiomers, may be separated using known methods, such as through the use of, for example, chiral salts and chiral chromatography.

In addition, it is recognized that one optical isomer, including a diastereomer and enantiomer, or a stereoisomer, may have favorable properties over the other. Thus, when disclosing and claiming the invention, when one racemic mixture is disclosed, it is clearly contemplated that both optical isomers, including diastereomers and enantiomers, or stereoisomers substantially free of the other are disclosed and claimed as well.

The syntheses of the compounds useful in the present invention are described in the art. Accordingly, the ordinarily skilled artisan will be able to prepare the compounds described herein. For further guidance, the syntheses of the present compounds are described in Yokoyama et al., U.S. Patent No. 5,401,772, assigned to Ciba-Geigy Corp., issued March 28, 1995; Yokoyama et al., EP 0,580,550, assigned to Ciba-Geigy Corp., published January 26, 1994; and Yokoyama et al., "Synthesis and Structure-Activity Relationships of Oxamic Acid and Acetic Acid Derivative Related to L-Thyronine", *Journal of Medicinal Chemistry*, Vol. 38, pp. 695 - 707 (1995).

For example, the following describes the synthesis of illustrative compounds utilized in the present invention.



Example 1

**1a. 2-isopropyl anisole:** Potassium hydroxide (5.6 g) is added to 13.4 mL acetone followed by 2-isopropylphenol (13.6 g). After the potassium hydroxide is dissolved, methyl iodide (14.2 g) is added. The reaction is refluxed for about 16 hours. 150 mL of water is added. This reaction is extracted 3 times with 100 mL diethyl ether. The organic layer is extracted twice with 100 mL 10% sodium hydroxide in water, once with 100 mL water, and once with 100 mL saturated ammonium chloride. After drying over magnesium sulfate, the organic solution is dried over MgSO<sub>4</sub>, filtered, and concentrated under reduced pressure. The material is fractionally distilled under reduced pressure to afford 1a.


**1b. Bis(3-isopropyl-4-methoxyphenyl)iodonium Tetrafluoroborate:** Acetic anhydride (7 mL) is cooled to  $-15^{\circ}\text{C}$  in a dry ice acetone bath. Fuming nitric acid (5.4 mL) is added dropwise. Iodine (2.5 g) is added in one piece followed by dropwise addition of 4.7 mL trifluoroacetic acid. After 20 minutes, the reaction is removed from the bath and stirred at room temperature for 30 minutes. After the iodine has dissolved, the reaction is sparged to remove nitrogen oxides and then concentrated under vacuum. The material is then taken up in 15 mL acetic anhydride and cooled to  $-10^{\circ}\text{C}$ . To this cooled solution is added dropwise a solution of 2-isopropyl anisole (**1a**; 7.43 g) in 35 mL acetic anhydride and 5 mL trifluoroacetic acid. The reaction is allowed to stand in a refrigerator for about 16 hours. After allowing the reaction to return to room temperature for 3 hours, the reaction is concentrated under high vacuum. The residue is taken up in 25 mL methanol, 25 mL 10% sodium bisulfite, and 188 mL 2M sodium tetrafluoroborate. The mixture is stirred vigorously for 30 minutes and the supernatant is decanted off. To the residue is added 200 mL hexane and it is stirred for an additional 30 minutes. The solid is collected, washed with hexane, and dried under vacuum to afford **1b**.

**1c. 2',6'-dimethyl-3-isopropyl-4-methoxy-4'-nitrodiphenyl ether:** Bis(3-isopropyl-4-methoxyphenyl)iodonium tetrafluoroborate (**1b**; 3 g) is weighed and taken up in 7.7 mL dichloromethane and 0.5 g copper bronze is added. The mixture is cooled in an ice water bath. A solution of 2,6-dimethyl-4-nitrophenol (0.65 g) and triethylamine (0.44 g) in 5.2 mL dichloromethane is added dropwise. The reaction is placed in the dark and stirred for 5 days. At this time, the reaction is filtered through celite and concentrated under reduced pressure. Purification of the product by chromatography on silica gel followed by crystallization from hexane:ethyl acetate affords **1c**.


**1d. Methyl *N*-[3,5-dimethyl-4-(4'-methoxy-3'-isopropylphenoxy)phenyl]oxamate:** 2',6'-dimethyl-3-isopropyl-4-methoxy-4'-nitrodiphenyl ether (**1c**; 0.51 g) is dissolved in 20 mL ethanol and 60 mg of 10% palladium on carbon is added. The reaction is hydrogenated for 3 hours, then filtered through celite and concentrated under reduced pressure. Dimethyl oxamate (3 g), is added to the residue and the reaction is heated to  $120^{\circ}\text{C}$  for 4 hours. The reaction mixture is concentrated under reduced pressure and purified by chromatography on silica gel to afford **1d**.

1e. *N*-[3,5-dimethyl-4-(4'-hydroxy-3'-isopropylphenoxy)phenyl]oxamate: Methyl *N*-[3,5-dimethyl-4-(4'-methoxy-3'-isopropylphenoxy)phenyl]oxamate (1d; 350 mg) is dissolved in 3.5 mL dichloromethane and cooled in a dry ice acetone bath. To this solution is dropwise added 2 mL boron tribromide (1 M in dichloromethane). The reaction is stirred about 16 hours and is allowed to reach ambient temperature. At this time, the reaction is poured onto 10 mL ice and water. To this mixture is added 10 mL ethyl acetate. The organic layer is separated and the aqueous phase is extracted twice with 10 mL ethyl acetate. The organic layers are combined, dried over magnesium sulfate, and concentrated under reduced pressure. The material is dissolved in 1:1 acetonitrile:water and loaded onto a column containing 5 g C<sub>18</sub>-derivatized silica gel and equilibrated with water. The product is eluted using a step gradient, 2 x 15 mL 25:75 acetonitrile:water, 1 x 15 mL 35:65 acetonitrile:water, 1 x 15 mL 40:60 acetonitrile:water, 2 x 15 mL 45:55 acetonitrile:water, and 1 x 15 mL 50:50 acetonitrile:water to afford 1e.


Example 2



The compound of Example 2 is synthesized as described in Yokoyama et al., "Synthesis and Structure-Activity Relationships of Oxamic Acid and Acetic Acid Derivative Related to L-Thyronine", *Journal of Medicinal Chemistry*, Vol. 38, pp. 695 - 707 (1995).

Example 3

The compound of Example 3 is synthesized as described in Yokoyama et al., U.S. Patent No. 5,401,772, assigned to Ciba-Geigy Corp., issued March 28, 1995.


Example 4

**4a. 2',6'-diiodo-3-isopropyl-4-methoxy-4'-nitrodiphenyl ether:** Bis(3-isopropyl-4-methoxyphenyl)iodonium tetrafluoroborate (prepared as described in Example 1b) (3 g) is weighed is taken up in 7.7 mL dichloromethane and 0.5 g copper bronze is added. The mixture is cooled in an ice water bath. A solution of 2,6-diiodo-4-nitrophenol (1.53 g) and triethyl amine (0.43 g) in 5 mL dichloromethane is added dropwise. The reaction is placed in the dark and stirred for 5 days. At this time, 25 mL methanol is added to the reaction and is filtered through a plug of silica gel. The silica gel is washed with an additional 5 mL of

methanol and the filtrate is concentrated under reduced pressure. Purification of the product by chromatography on silica gel affords 4a.

**4b. Methyl *N*-[3,5-diiodo-4-(4'-methoxy-3'-isopropylphenoxy)phenyl]oxamate:** 2',6'-diiodo-3-isopropyl-4-methoxy-4'-nitrodiphenyl ether (4a) (0.62 g) is dissolved in 20 mL ethanol and 2 mL *N,N*-dimethylformamide and 40 mg of 10% palladium on carbon is added. The reaction is hydrogenated for 4 hours, then filtered through celite and concentrated under reduced pressure. Dimethyl oxamate (3g) is added to the residue and the reaction is heated to 120 °C for 3 hours. The reaction mixture is concentrated under reduced pressure and purified by chromatography on silica gel to afford 4b.

**4c. *N*-[3,5-diiodo-4-(4'-hydroxy-3'-isopropylphenoxy)phenyl]oxamate:** Methyl *N*-[3,5-diiodo-4-(4'-methoxy-3'-isopropylphenoxy)phenyl]oxamate (4b) (62 mg) is dissolved in 1 mL dichloromethane and cooled in a dry ice acetone bath. To this solution is added 0.2 mL boron tribromide (1 M in dichloromethane). The reaction is stirred for about 16 hours and allowed to reach room temperature. The reaction is poured onto 10 mL ice and water. To this mixture is added 10 mL ethyl acetate. The organic layer is separated and the aqueous phase is extracted a second time with 10 mL ethyl acetate. The organic layers are combined, washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. Purification of the product by chromatography on silica gel affords 4c.



**5. *N*-[3,5-dimethyl-4-(4'-methoxy-3'-isopropylphenoxy)phenyl]oxamate:** Methyl *N*-[3,5-dimethyl-4-(4'-methoxy-3'-isopropylphenoxy)phenyl]oxamate (prepared as described in Example 1d) (750 mg) is dissolved in 12 mL methanol and 2.2 mL 1 N sodium hydroxide is added. The sample is reacted overnight. The mixture is concentrated under reduced pressure. To the mixture is added 25 mL water and 15 mL ethyl acetate, followed by 5 mL 1

N HCl. The mixture is transferred to a separatory funnel and an additional 35 mL ethyl acetate is added. The organic layer is separated and the aqueous phase is extracted once with 15 mL ethyl acetate and once with 10 mL ethyl acetate. The organic layers are combined, washed with 25 mL saturated sodium chloride, and dried over magnesium sulfate. The solution is filtered and concentrated under reduced pressure to afford 5.

#### Use of the Present Compounds

The methods of the present invention are performed by administering to a mammal (preferably a human) a compound having a structure as described herein and, preferably, a pharmaceutically-acceptable or cosmetically-acceptable carrier.

The compounds herein may be used for the treatment of such conditions as treating hair loss in mammals, including arresting and / or reversing hair loss and promoting hair growth. Such conditions may manifest themselves in, for example, alopecia, including male pattern baldness and female pattern baldness.

Preferably the compounds of the present invention are, as defined herein, cardiac-sparing.

Preferably, in the methods of the present invention, the compounds are formulated into pharmaceutical or cosmetic compositions for use in treatment or prophylaxis of conditions such as the foregoing. Standard pharmaceutical formulation techniques are used, such as those disclosed in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA. (1990).

Typically, from about 5 mg to about 3000 mg, more preferably from about 5 mg to about 1000 mg, more preferably from about 10 mg to about 100 mg, of a compound having a structure as described herein is administered per day for systemic administration. It is understood that these dosage ranges are by way of example only, and that daily administration can be adjusted depending on various factors. The specific dosage of the compound to be administered, as well as the duration of treatment, and whether the treatment is topical or systemic are interdependent. The dosage and treatment regimen will also depend upon such factors as the specific compound used, the treatment indication, the efficacy of the compound, the personal attributes of the subject (such as, for example, weight, age, sex, and medical condition of the subject), compliance with the treatment regimen, and the presence and severity of any side effects of the treatment.

According to the present invention, the subject compounds are co-administered with a pharmaceutically-acceptable or cosmetically-acceptable carrier (herein collectively described as "carrier"). The term "carrier", as used herein, means one or more compatible solid or liquid filler diluents or encapsulating substances which are suitable for administration to a mammal. The term "compatible", as used herein, means that the components of the composition are capable of being commingled with a compound of the present invention, and with each other, in a manner such that there is no interaction which would substantially reduce the efficacy of the composition under ordinary use situations. Carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the animal, preferably mammal (most preferably human), being treated. The carrier can itself be inert or it can possess pharmaceutical and / or cosmetic benefits of its own.

The compositions of this invention may be in any of a variety of forms, suitable (for example) for oral, rectal, topical, nasal, ocular or parenteral administration. Of these, topical and / or oral administration are especially preferred with topical being most preferred. Depending upon the particular route of administration desired, a variety of carriers well-known in the art may be used. These include solid or liquid fillers, diluents, hydrotropes, surface-active agents, and encapsulating substances. Optional pharmaceutically-active or cosmetically-active materials may be included which do not substantially interfere with the activity of the compound of the present invention. The amount of carrier employed in conjunction with the compound is sufficient to provide a practical quantity of material for administration per unit dose of the compound. Techniques and compositions for making dosage forms useful in the methods of this invention are described in the following references: Modern Pharmaceutics, Chapters 9 and 10, Banker & Rhodes, eds. (1979); Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms, 2<sup>nd</sup> Ed., (1976).

Some examples of substances which can serve as carriers or components thereof are sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, and methyl cellulose; powdered tragacanth; malt; gelatin; talc; solid lubricants, such as stearic acid and magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerine, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers, such as the

TWEENS; wetting agents, such sodium lauryl sulfate; coloring agents; flavoring agents; tableting agents, stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline; and phosphate buffer solutions.

The choice of a carrier to be used in conjunction with the subject compound is typically determined by the way the compound is to be administered.

In particular, carriers for systemic administration include sugars, starches, cellulose and its derivatives, malt, gelatin, talc, calcium sulfate, vegetable oils, synthetic oils, polyols, alginic acid, phosphate buffer solutions, emulsifiers, isotonic saline, and pyrogen-free water. Preferred carriers for parenteral administration include propylene glycol, ethyl oleate, pyrrolidone, ethanol, and sesame oil. Preferably, the carrier, in compositions for parenteral administration, comprises at least about 90% by weight of the total composition.

Various oral dosage forms can be used, including such solid forms as tablets, capsules, granules and bulk powders. These oral forms comprise a safe and effective amount, usually at least about 5%, and preferably from about 25% to about 50%, of a compound used in the present invention. Tablets can be compressed, tablet triturates, enteric-coated, sugar-coated, film-coated, or multiple-compressed, containing suitable binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. Liquid oral dosage forms include aqueous solutions, emulsions, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules, and effervescent preparations reconstituted from effervescent granules, containing suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, melting agents, coloring agents and flavoring agents.

The carriers suitable for the preparation of unit dosage forms for oral administration are well-known in the art. Tablets typically comprise conventional pharmaceutically-compatible adjuvants as inert diluents, such as calcium carbonate, sodium carbonate, mannitol, lactose and cellulose; binders such as starch, gelatin and sucrose; disintegrants such as starch, alginic acid and croscarmellose; lubricants such as magnesium stearate, stearic acid and talc. Glidants such as silicon dioxide can be used to improve flow characteristics of the powder mixture. Coloring agents, such as the FD&C dyes, can be added for appearance. Sweeteners and flavoring agents, such as aspartame, saccharin, menthol, peppermint, and fruit flavors, are useful adjuvants for chewable tablets. Capsules (including time release and sustained release formulations) typically comprise one or more solid diluents disclosed above. The selection of carrier components depends on secondary considerations like taste,



cost, and shelf stability, which are not critical for the purposes of the subject invention, and can be readily made by a person skilled in the art.

Orally administered compositions also include liquid solutions, emulsions, suspensions, powders, granules, elixirs, tinctures, syrups, and the like. The carriers suitable for preparation of such compositions are well known in the art. Typical components of carriers for syrups, elixirs, emulsions and suspensions include ethanol, glycerol, propylene glycol, polyethylene glycol, liquid sucrose, sorbitol and water. For a suspension, typical suspending agents include methyl cellulose, sodium carboxymethyl cellulose, AVICEL RC-591, tragacanth and sodium alginate; typical wetting agents include lecithin and polysorbate 80; and typical preservatives include methyl paraben and sodium benzoate. Peroral liquid compositions may also contain one or more components such as sweeteners, flavoring agents and colorants disclosed above.

Such compositions may also be coated by conventional methods, typically with pH or time-dependent coatings, such that the subject compound is released in the gastrointestinal tract in the vicinity of the desired topical application, or at various times to extend the desired action. Such dosage forms typically include, but are not limited to, one or more of cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropyl methyl cellulose phthalate, ethyl cellulose, Eudragit coatings, waxes and shellac.

Other compositions useful for attaining systemic delivery of the subject compounds include sublingual, buccal and nasal dosage forms. Such compositions typically comprise one or more of soluble filler substances such as sucrose, sorbitol and mannitol; and binders such as acacia, microcrystalline cellulose, carboxymethyl cellulose and hydroxypropyl methyl cellulose. Glidants, lubricants, sweeteners, colorants, antioxidants and flavoring agents disclosed above may also be included.

The compounds of the present invention may also be topically administered. The carrier of the topical composition preferably aids penetration of the present compounds into the skin to reach the environment of the hair follicle. Topical compositions of the present invention may be in any form including, for example, solutions, oils, creams, ointments, gels, lotions, shampoos, leave-on and rinse-out hair conditioners, milks, cleansers, moisturizers, sprays, skin patches, and the like.

Topical compositions containing the active compound can be admixed with a variety of carrier materials well known in the art, such as, for example, water, alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, propylene glycol, PPG-2 myristyl propionate, and the like.

Other materials suitable for use in topical carriers include, for example, emollients, solvents, humectants, thickeners and powders. Examples of each of these types of materials, which can be used singly or as mixtures of one or more materials, are as follows:

Emollients, such as stearyl alcohol, glyceryl monoricinoleate, glyceryl monostearate, propane-1,2-diol, butane-1,3-diol, mink oil, cetyl alcohol, *iso*-propyl isostearate, stearic acid, *iso*-butyl palmitate, isocetyl stearate, oleyl alcohol, isopropyl laurate, hexyl laurate, decyl oleate, octadecan-2-ol, isocetyl alcohol, cetyl palmitate, dimethylpolysiloxane, di-*n*-butyl sebacate, *iso*-propyl myristate, *iso*-propyl palmitate, *iso*-propyl stearate, butyl stearate, polyethylene glycol, triethylene glycol, lanolin, sesame oil, coconut oil, arachis oil, castor oil, acetylated lanolin alcohols, petroleum, mineral oil, butyl myristate, isostearic acid, palmitic acid, isopropyl linoleate, lauryl lactate, myristyl lactate, decyl oleate, and myristyl myristate; propellants, such as propane, butane, *iso*-butane, dimethyl ether, carbon dioxide, and nitrous oxide; solvents, such as ethyl alcohol, methylene chloride, *iso*-propanol, castor oil, ethylene glycol monoethyl ether, diethylene glycol monobutyl ether, diethylene glycol monoethyl ether, dimethyl sulphoxide, dimethyl formamide, tetrahydrofuran; humectants, such as glycerin, sorbitol, sodium 2-pyrrolidone-5-carboxylate, soluble collagen, dibutyl phthalate, and gelatin; and powders, such as chalk, talc, fullers earth, kaolin, starch, gums, colloidal silicon dioxide, sodium polyacrylate, tetra alkyl ammonium smectites, trialkyl aryl ammonium smectites, chemically modified magnesium aluminium silicate, organically modified montmorillonite clay, hydrated aluminium silicate, fumed silica, carboxyvinyl polymer, sodium carboxymethyl cellulose, and ethylene glycol monostearate.

The compounds used in the present invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines. A preferred formulation for topical delivery of the present compounds utilizes liposomes such as described in Dowton et al., "Influence of Liposomal Composition on Topical Delivery of Encapsulated Cyclosporin A: I. An *in vitro* Study Using Hairless Mouse Skin", *S.T.P. Pharma Sciences*, Vol. 3, pp. 404 - 407 (1993); Wallach and Philippot, "New Type of Lipid Vesicle: Novasome<sup>®</sup>", Liposome Technology, Vol. 1, pp. 141 - 156 (1993); Wallach, U.S. Patent No. 4,911,928, assigned to Micro-Pak, Inc., issued March 27, 1990; and Weiner et al., U.S. Patent No. 5,834,014, assigned to The University of Michigan and Micro-Pak, Inc., issued November 10, 1998 (with respect to Weiner et al., with a compound as described herein administered in lieu of, or in addition to, minoxidil).

The compounds of the present invention may also be administered by iontophoresis. See, e.g., internet site [www.unipr.it/arpa/dipfarm/erasmus/erasm14.html](http://www.unipr.it/arpa/dipfarm/erasmus/erasm14.html); Banga et al., "Hydrogel-based Iontotherapeutic Delivery Devices for Transdermal Delivery of Peptide/Protein Drugs", *Pharm. Res.*, Vol. 10 (5), pp. 697-702 (1993); Ferry, "Theoretical Model of Iontophoresis Utilized in Transdermal Drug Delivery", *Pharmaceutical Acta Helvetiae*, Vol 70, pp. 279-287 (1995); Gangarosa et al., "Modern Iontophoresis for Local Drug Delivery", *Int. J. Pharm.*, Vol. 123, pp. 159-171 (1995); Green et al., "Iontophoretic Delivery of a Series of Tripeptides Across the Skin *in vitro*", *Pharm. Res.*, Vol 8, pp. 1121-1127 (1991); Jadoul et al., "Quantification and Localization of Fentanyl and TRH Delivered by Iontophoresis in the Skin", *Int. J. Pharm.*, Vol. 120, pp. 221-8 (1995); O'Brien et al., "An Updated Review of its Antiviral Activity, Pharmacokinetic Properties and Therapeutic Efficacy", *Drugs*, Vol. 37, pp. 233-309 (1989); Parry et al., "Acyclovir Bioavailability in Human Skin", *J. Invest. Dermatol.*, Vol. 98 (6), pp. 856-63 (1992); Santi et al., "Drug Reservoir Composition and Transport of Salmon Calcitonin in Transdermal Iontophoresis", *Pharm. Res.*, Vol 14 (1), pp. 63-66 (1997); Santi et al., "Reverse Iontophoresis - Parameters Determining Electroosmotic Flow: I. pH and Ionic Strength", *J. Control. Release*, Vol. 38, pp. 159-165 (1996); Santi et al., "Reverse Iontophoresis - Parameters Determining Electroosmotic Flow: II. Electrode Chamber Formulation", *J. Control. Release*, Vol. 42, pp. 29-36 (1996); Rao et al., "Reverse Iontophoresis: Noninvasive Glucose Monitoring *in vivo* in Humans", *Pharm. Res.*, Vol. 12 (12), pp. 1869-1873 (1995); Thysman et al., "Human Calcitonin Delivery in Rats by Iontophoresis", *J. Pharm. Pharmacol.*, Vol. 46, pp. 725-730 (1994); and Volpato et al., "Iontophoresis Enhances the Transport of Acyclovir through Nude Mouse Skin by Electrorepulsion and Electroosmosis", *Pharm. Res.*, Vol. 12 (11), pp. 1623-1627 (1995).

The compositions used in the present invention may also optionally comprise an activity enhancer. The activity enhancer can be chosen from a wide variety of molecules which can function in different ways to enhance hair growth effects of a compound of the present invention. Particular classes of activity enhancers include other hair growth stimulants and penetration enhancers.

Non-limiting examples of other hair growth stimulants which may be used in the compositions herein, including both systemic and topical compositions, include, for example, benzalkonium chloride, benzethonium chloride, phenol, estradiol, diphenhydramine hydrochloride, chlorpheniramine maleate, chlorophyllin derivatives, cholesterol, salicylic acid, cysteine, methionine, red pepper tincture, benzyl nicotinate, D,L - menthol, peppermint

oil, calcium pantothenate, panthenol, castor oil, hinokitiol, prednisolone, resorcinol, monosaccharides and esterified monosaccharides, chemical activators of protein kinase C enzymes, glycosaminoglycan chain cellular uptake inhibitors, inhibitors of glycosidase activity, glycosaminoglycanase inhibitors, esters of pyroglutamic acid, hexosaccharic acids or acylated hexosaccharic acids, aryl-substituted ethylenes, N-acylated amino acids, and, of course, minoxidil or finasteride. The most preferred activity enhancers are minoxidil and finasteride, most preferably minoxidil.

Non-limiting examples of penetration enhancers which may be used in the compositions herein include, for example, 2-methyl propan-2-ol, propan-2-ol, ethyl-2-hydroxypropanoate, hexan-2,5-diol, POE(2) ethyl ether, di(2-hydroxypropyl) ether, pentan-2,4-diol, acetone, POE(2) methyl ether, 2-hydroxypropionic acid, 2-hydroxyoctanoic acid, propan-1-ol, 1,4-dioxane, tetrahydrofuran, butan-1,4-diol, propylene glycol dipelargonate, polyoxypropylene 15 stearyl ether, octyl alcohol, POE ester of oleyl alcohol, oleyl alcohol, lauryl alcohol, dioctyl adipate, dicapryl adipate, di-isopropyl adipate, di-isopropyl sebacate, dibutyl sebacate, diethyl sebacate, dimethyl sebacate, dioctyl sebacate, dibutyl suberate, dioctyl azelate, dibenzyl sebacate, dibutyl phthalate, dibutyl azelate, ethyl myristate, dimethyl azelate, butyl myristate, dibutyl succinate, didecyl phthalate, decyl oleate, ethyl caproate, ethyl salicylate, *iso*-propyl palmitate, ethyl laurate, 2-ethyl-hexyl pelargonate, *iso*-propyl isostearate, butyl laurate, benzyl benzoate, butyl benzoate, hexyl laurate, ethyl caprate, ethyl caprylate, butyl stearate, benzyl salicylate, 2-hydroxypropanoic acid, 2-hydroxyoctanoic acid, dimethyl sulphoxide, N,N-dimethyl acetamide, N,N-dimethyl formamide, 2-pyrrolidone, 1-methyl-2-pyrrolidone, 5-methyl-2-pyrrolidone, 1,5-dimethyl-2-pyrrolidone, 1-ethyl-2-pyrrolidone, phosphine oxides, sugar esters, tetrahydrofurfural alcohol, urea, diethyl-*m*-toluamide, and, 1-dodecylazacycloheptan-2-one.

In all of the foregoing, of course, the compounds used in the present methods can be administered alone or as mixtures, and the compositions may further include additional drugs or excipients as appropriate for the indication.

The present invention further relates to kits comprising a compound and / or composition herein and information and / or instructions by words, pictures, and / or the like, that use of the kit will provide treatment for hair loss in mammals (particularly humans) including, for example, arresting and / or reversing hair loss and / or promoting hair growth. In addition or in the alternative, the kit may comprise a compound and / or composition herein and information and / or instructions regarding methods of application of the compound and / or composition, preferably with the benefit of treating hair loss in mammals.

Examples of Composition Administration

The following examples do not limit the invention, but provide guidance to the skilled artisan to perform the methods of the present invention. In each example, a compound other than the one mentioned may be substituted in the example by another having a structure as described herein with similar results.

Example A

A composition for topical administration is made, comprising:

<u>Component</u>	<u>Amount</u>
Compound of Example 1	5 %
Ethanol	57 %
Propylene Glycol	19 %
Dimethyl Isosorbide	19 %

A human male subject suffering from male pattern baldness is treated by a method of this invention. Specifically, for 6 weeks, the above composition is daily administered topically to the subject.

Example B

A composition for topical administration is made according to the method of Dowton et al., "Influence of Liposomal Composition on Topical Delivery of Encapsulated Cyclosporin A: I. An *in vitro* Study Using Hairless Mouse Skin", S.T.P. Pharma Sciences, Vol. 3, pp. 404 - 407 (1993), using 3,5-Example 1 in lieu of cyclosporin A and using the Novasome 1 for the non-ionic liposomal formulation.

A human male subject suffering from male pattern baldness is treated each day with the above composition. Specifically, for 6 weeks, the above composition is administered topically to the subject.

Example C

A shampoo is made, comprising:


	Ex. C-1	Ex. C-2	Ex. C-3	Ex. C-4
<u>Component</u>				
Ammonium Lauryl Sulfate	11.5 %	11.5 %	9.5 %	7.5 %
Ammonium Laureth Sulfate	4 %	3 %	2 %	2 %
Cocamide MEA	2 %	2 %	2 %	2 %

Ethylene Glycol Distearate	2 %	2 %	2 %	2 %
Cetyl Alcohol	2 %	2 %	2 %	2 %
Stearyl Alcohol	1.2 %	1.2 %	1.2 %	1.2 %
Glycerin	1 %	1 %	1 %	1 %
Polyquaternium 10	0.5 %	0.25 %	-	-
Polyquaternium 24	-	-	0.5 %	0.25 %
Sodium Chloride	0.1 %	0.1 %	0.1 %	0.1 %
Sucrose Polyesters of Cottonate Fatty Acid	3 %	3 %	-	-
Sucrose Polyesters of Behenate Fatty Acid	2 %	3 %	-	-
Polydimethyl Siloxane	-	-	3 %	2 %
Cocaminopropyl Betaine	-	1 %	3 %	3 %
Lauryl Dimethyl Amine Oxide	1.5 %	1.5 %	1.5 %	1.5 %
Decyl Polyglucose	-	-	1 %	1 %
DMDM Hydantoin	0.15 %	0.15 %	0.15 %	0.15 %
Compound of Example 1	4 %	-	3 %	-
Compound of Example 2	-	5 %	-	-
Compound of Example 3	-	-	-	4 %
Minoxidil	-	-	3 %	2 %
Phenoxyethanol	0.5 %	0.5 %	0.5 %	0.5 %
Fragrance	0.5 %	0.5 %	0.5 %	0.5 %
Water	q.s.	q.s.	q.s.	q.s.

A human subject suffering from male pattern baldness is treated by a method of this invention. Specifically, for 12 weeks, the above shampoo is used daily by the subject.

What is claimed is:

1. A method of treating hair loss comprising administering a composition comprising a cardiac-sparing compound characterized by the structure:




and pharmaceutically acceptable salts, hydrates, and biohydrolyzable amides, esters, and imides thereof, wherein:

R is selected from the group consisting of hydrogen, hydroxy, esterified hydroxy, and etherified hydroxy;

R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> are each, independently, selected from the group consisting of hydrogen, halogen, trifluoromethyl and lower alkyl;

R<sub>3</sub> is selected from the group consisting of halogen, trifluoromethyl, lower alkyl, aryl, aryl-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, and:



R<sub>8</sub> is selected from the group consisting of hydrogen, lower alkyl, aryl, cycloalkyl, aryl-lower alkyl, and cycloalkyl-lower alkyl;

R<sub>9</sub> is selected from the group consisting of hydroxy and acyloxy;

R<sub>10</sub> is selected from the group consisting of hydrogen and lower alkyl; with the proviso that R<sub>9</sub> and R<sub>10</sub> collectively optionally represent oxo;

W is selected from the group consisting of -O- and -S-;

X is selected from the group consisting of -NR<sub>7</sub>, S, and O;

R<sub>5</sub> is selected from the group consisting of hydrogen, lower alkyl, and aryl-lower alkyl and R<sub>6</sub> is hydrogen; with the proviso that when X is -NR<sub>7</sub>, R<sub>5</sub> and R<sub>6</sub> together are optionally oxo;

R<sub>7</sub> is selected from the group consisting of hydrogen and lower alkyl; and

Z is selected from the group consisting of carboxyl and carboxyl derivatized as a pharmaceutically acceptable ester or amide.

2. A method according to claim 1 wherein X is -NR<sub>7</sub> and R<sub>5</sub> and R<sub>6</sub> together are oxo.

3. A method according to any of claims 1 and 2 wherein:

R substitutes on the 4' position;

R<sub>1</sub> substitutes on the 3 position;

R<sub>2</sub> substitutes on the 5 position;

R<sub>3</sub> substitutes on the 3' position; and

R<sub>4</sub> substitutes on the 5' position.


4. A method according any of claims 1 and 2 wherein:

W is -O-;

R is selected from the group consisting of hydroxy, esterified hydroxy, and etherified hydroxy;

R<sub>4</sub> is hydrogen;


R<sub>3</sub> is selected from the group consisting of lower alkyl, aryl-lower alkyl, cycloalkyl-lower alkyl and



Z is selected from the group consisting of carboxyl and carboxyl derivatized as a pharmaceutically acceptable ester.

5. A method according to claim 4 wherein the compound is characterized by the structure:





wherein R<sub>1</sub> and R<sub>2</sub> are each, independently, selected from the group consisting of hydrogen, halogen, trifluoromethyl, and C<sub>1</sub> - C<sub>3</sub> alkyl.

6. A method according claim 5 wherein R<sub>7</sub> is selected from the group consisting of hydrogen and C<sub>1</sub> - C<sub>3</sub> alkyl.
7. A method according to any of claims 5 and 6 wherein R is hydroxy.
8. A method according to any of claims 5, 6, and 7 wherein R<sub>3</sub> is selected from the group consisting of isopropyl, benzyl, benzyl substituted with halogen, and trifluoromethyl.
9. A method according to any of claims 5, 6, 7, and 8 wherein Z is carboxyl.
10. A method according to any of the preceding claims wherein the administration is topical.

## INTERNATIONAL SEARCH REPORT

Internat al Application No

PCT/US 00/05253

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K7/06 A61P17/14

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 580 550 A (CIBA GEIGY AG) 26 January 1994 (1994-01-26) page 2, line 2 - line 15; claims 1,18 ---	1,5
A	CHEMICAL ABSTRACTS, vol. 82, no. 15, 14 April 1975 (1975-04-14) Columbus, Ohio, US; abstract no. 93567w, P.A. HALE ET AL: "Effects of epilation and hormones on the activity of rat hair follicles" page 81; XP002141613 abstract & J. EXP. ZOOL., vol. 191, no. 1, 1975, pages 49-62, --- -/--	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

7 July 2000

Date of mailing of the international search report

24/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Voyiazoglou, D

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05253

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 284 971 A (WALKER KEITH A ET AL) 8 February 1994 (1994-02-08) cited in the application column 1 -column 2 ---	1,5
A	US 5 061 798 A (EMMETT JOHN COLIN ET AL) 29 October 1991 (1991-10-29) cited in the application column 16, line 55 -column 17, line 67; claim 1 ---	1,5
A	W. J. CUNLIFFE ET AL : "Alopecia areata, thyroid disease and autoimmunity" BRITISH JOURNAL OF DERMATOLOGY, vol. 81, no. 12, 1969, pages 877-881, XP000923239 page 67 ---	1
A	EP 0 123 528 A (MORTIMER CHRISTOPHER H DR) 31 October 1984 (1984-10-31) cited in the application claim 1 ---	1
A	EP 0 749 752 A (CIRD GALDERMA) 27 December 1996 (1996-12-27) claims 1,3 ---	1
A	US 4 061 791 A (HALL CHARLES M ET AL) 6 December 1977 (1977-12-06) claims 1,17 ---	1,5
A	WO 94 27586 A (HANDELMAN JOSEPH H ;AHLUWALIA GURPREET S (US); SHANDER DOUGLAS (US) 8 December 1994 (1994-12-08) claims 1,2 ---	1,5
A	EP 0 898 956 A (OREAL) 3 March 1999 (1999-03-03) page 4, line 30 -page 5, line 35; claim 1 ---	1,5
A	US 4 939 171 A (MOELLER HINRICH ET AL) 3 July 1990 (1990-07-03) claims 1,9 ---	1
A	EP 0 656 350 A (SQUIBB BRISTOL MYERS CO) 7 June 1995 (1995-06-07) page 8, line 3 - line 8; example 15 ---	1,5
	--- -/--	

# INTERNATIONAL SEARCH REPORT

Intern: 31 Application No

PCT/US 00/05253

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	<p>CHEMICAL ABSTRACTS, vol. 131, no. 17,  25 October 1999 (1999-10-25)  Columbus, Ohio, US;  abstract no. 223774e,  V. K. SHARMA ET AL : "Evaluation of  thyroid function in North Indians with  alopecia areata: response to intravenous  injection of 100 microgram thyrotropin  releasing hormone"  page 125;  XP002141564  abstract  &amp; J. DERMATOL.,  vol. 26, no. 6, 1999, pages 339-342,  -----</p>	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/05253

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0580550 A	26-01-1994	AT 159515 T AU 4208193 A CA 2100817 A DE 69314718 D DE 69314718 T DK 580550 T ES 2108855 T FI 933260 A GR 3025517 T HU 64512 A,B JP 6172275 A NO 932614 A NZ 248181 A US 5401772 A US 5569674 A US 5654468 A ZA 9305196 A	15-11-1997 27-01-1994 22-01-1994 27-11-1997 26-02-1998 02-02-1998 01-01-1998 22-01-1994 27-02-1998 28-01-1994 21-06-1994 24-01-1994 27-11-1995 28-03-1995 29-10-1996 05-08-1997 07-07-1994
US 5284971 A	08-02-1994	NONE	
US 5061798 A	29-10-1991	AT 61581 T AU 577917 B AU 5221986 A CA 1319148 A CN 1010310 B DE 3678000 D DK 18586 A,B, EP 0188351 A ES 551005 D ES 8800157 A FI 860229 A GR 860122 A HU 40401 A,B IE 58867 B IL 77605 A JP 7103070 B JP 61167643 A KR 9311302 B NO 860159 A NZ 214843 A PH 22691 A PT 81844 A,B US 4826876 A US 4910305 A US 4766121 A ZA 8600319 A ZW 886 A	15-03-1991 06-10-1988 24-07-1986 15-06-1993 07-11-1990 18-04-1991 19-07-1986 23-07-1986 01-11-1987 01-01-1988 19-07-1986 19-05-1986 28-12-1986 17-11-1993 09-02-1990 08-11-1995 29-07-1986 29-11-1993 21-07-1986 24-02-1989 14-11-1988 01-02-1986 02-05-1989 20-03-1990 23-08-1988 27-08-1986 13-05-1987
EP 0123528 A	31-10-1984	GB 2138286 A AT 29382 T AU 566833 B AU 2828484 A CA 1226524 A DE 3465879 D DK 607284 A WO 8404038 A GR 81975 A JP 60501057 T	24-10-1984 15-09-1987 29-10-1987 07-11-1984 08-09-1987 15-10-1987 18-12-1984 25-10-1984 12-12-1984 11-07-1985

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat'l Application No

PCT/US 00/05253

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0123528 A		NO 845096 A NZ 207923 A PT 78456 A ZA 8402986 A	18-12-1984 28-07-1988 01-05-1984 26-06-1985
EP 0749752 A	27-12-1996	FR 2735367 A AU 688216 B AU 5470396 A CA 2179426 A JP 2804014 B JP 9002972 A US 6004987 A	20-12-1996 05-03-1998 16-01-1997 20-12-1996 24-09-1998 07-01-1997 21-12-1999
US 4061791 A	06-12-1977	US 4128570 A US 4159341 A US 4156012 A	05-12-1978 26-06-1979 22-05-1979
WO 9427586 A	08-12-1994	AU 677346 B AU 7018194 A CA 2163535 A EP 0700288 A JP 8510734 T ZA 9403611 A	17-04-1997 20-12-1994 08-12-1994 13-03-1996 12-11-1996 25-01-1995
EP 0898956 A	03-03-1999	FR 2767689 A JP 3023088 B JP 11116425 A	05-03-1999 21-03-2000 27-04-1999
US 4939171 A	03-07-1990	DE 3738406 A AT 77047 T DE 3871904 A EP 0315913 A GR 3005612 T JP 1153617 A	24-05-1989 15-06-1992 16-07-1992 17-05-1989 07-06-1993 15-06-1989
EP 0656350 A	07-06-1995	US 5547966 A AU 690133 B AU 7446394 A CA 2132771 A JP 7188151 A	20-08-1996 23-04-1998 27-04-1995 08-04-1995 25-07-1995